

業績

Sawada Y, Hangai M, Ishikawa M, Yoshitomi T. Association of myopic deformation of optic disc with visual field progression in paired eyes with open-angle glaucoma. *PLoS One*. 2017;12(1): e0170733.

Kumagai K, Hangai M, Furukawa M, Suetsugu T, Ogino N. Centripetal movement of the capillaries in the central macular region after internal limiting membrane peeling. *Retin Cases Brief Rep*, in press.

Kumagai K, Uemura A, Hangai M, Suetsugu T, Ogino N. Effect of vitreomacular separation on macular thickness determined by spectral-domain optical coherence tomography. *Jpn J Ophthalmol*. 2017;61(1):84-91.

Tanaka D, Nakanishi H, Hangai M, Akagi T, Morooka S, Ikeda HO, Yoshimura N. Influence of high myopia on outcomes of trabeculectomy with mitomycin C in patients with primary open-angle glaucoma. *Jpn J Ophthalmol*. 2016;60(6):446-453.

Kumagai K, Hangai M, Furukawa M, Suetsugu T, Ogino N. Effect of vitreomacular adhesion and vitreous gel on age-related reduction of macular thickness: a retrospective observational study. *BMJ Open*. 2016;6(9):e012972.

Sawada Y, Hangai M, Ishikawa M, Yoshitomi T. Association of Myopic Optic Disc Deformation with Visual Field Defects in Paired Eyes with Open-Angle Glaucoma: A Cross-Sectional Study. *PLoS One*. 2016;11(8):e0161961.

Araie M, Murata H, Iwase A, Hangai M, Sugiyama K, Yoshimura N. Differences in Relationship Between Macular Inner Retinal Layer Thickness and Retinal Sensitivity in Eyes With Early and Progressed Glaucoma. *Invest Ophthalmol Vis Sci*. 2016;57(4):1588-94.

Kumagai K, Hangai M. Author Response: progressive thinning of regional macular thickness after epiretinal membrane surgery. *Invest Ophthalmol Vis Sci*. 2016 Mar;57(3):1236-7.

Higashide T, Ohkubo S, Hangai M, Ito Y, Shimada N, Ohno-Matsui K, Terasaki H, Sugiyama K, Chew P, Li KK, Yoshimura N. Influence of Clinical Factors and Magnification Correction on Normal Thickness Profiles of Macular Retinal Layers Using Optical Coherence Tomography. *PLoS One*. 2016;11(1):e0147782.

Hasegawa T, Akagi T, Hangai M, Yamada H, Suda K, Kimura Y, Nakanishi H, Ikeda HO, Yoshimura N. Structural dissociation of optic disc margin components with optic disc tilting: a spectral domain optical coherence tomography study. *Graefes Arch Clin Exp Ophthalmol*. 2016 Feb;254(2):343-9.

Nakanishi H, Akagi T, Hangai M, Kimura Y, Suda K, Hasegawa T, Yamada H, Yoshikawa M, Morooka S, Ikeda HO, Yoshimura N. Effect of axial length on macular ganglion cell complex thickness and on early glaucoma diagnosis by spectral-domain optical coherence tomography. *J Glaucoma*. 2016 May;25(5):e481-90.

Kimura Y, Akagi T, Miyake M, Yamashiro K, Yoshikawa M, Yamada H, Hasegawa T, Suda K, Nakanishi H, Ohashi-Ikeda H, Gotoh N, Hangai M, Moriyama M, Ohno-Matsui K, Yoshimura N. Association between the CDKN2B-AS1 gene and primary open angle glaucoma with high myopia in japanese patients. *Ophthalmic Genet*. 2016 Jun;37(2):242-4.

Kumagai K, Hangai M. Author response: progressive thinning of regional macular thickness after epiretinal membrane surgery. *Invest Ophthalmol Vis Sci*. 2016;57(3):1236-7.

Higashide T, Ohkubo S, Hangai M, Ito Y, Shimada N, Ohno-Matsui K, Terasaki H, Sugiyama K, Chew P, Li KK, Yoshimura N. Influence of clinical factors and magnification correction on normal thickness profiles of macular retinal layers using optical coherence tomography. *PLoS One*. 2016;11(1):

Hasegawa T, Akagi T, Hangai M, Yamada H, Suda K, Kimura Y, Nakanishi H, Ikeda HO, Yoshimura N. Structural dissociation of optic disc margin components with optic disc tilting: a spectral domain optical coherence tomography study. *Graefes Arch Clin Exp Ophthalmol*. 2016;254(2):343-9.

Nakanishi H, Akagi T, Hangai M, Kimura Y, Suda K, Hasegawa T, Yamada H, Yoshikawa M, Morooka S, Ikeda HO, Yoshimura N. Effect of axial length on macular ganglion cell complex thickness and on early glaucoma diagnosis by spectral-domain optical coherence tomography. *Glaucoma*. 2015 in press.

Kumagai K, Hangai M, Ogino N. Progressive thinning of regional macular thickness after epiretinal membrane surgery. *Invest Ophthalmol Vis Sci*. 2015;56(12):7236-42.

Hosoda Y, Ooto S, Hangai M, Oishi A, Yoshimura N. Foveal photoreceptor deformation as a significant predictor of postoperative visual outcome in idiopathic epiretinal membrane surgery. *Invest Ophthalmol Vis Sci*. 2015;56(11):6387-93.

Kimura Y, Akagi T, Miyake M, Yamashiro K, Yoshikawa M, Yamada H, Hasegawa T, Suda K, Nakanishi H, Ohashi-Ikeda H, Gotoh N, Hangai M, Moriyama M, Ohno-Matsui K, Yoshimura N. Association between the *cdkn2b-as1* gene and primary open angle glaucoma with high myopia in Japanese patients. *Ophthalmic Genet*. 2015 in press.

Sawada Y, Hangai M, Murata K, Ishikawa M, Yoshitomi T. Lamina Cribrosa Depth Variation Measured by Spectral-Domain Optical Coherence Tomography Within and Between Four Glaucomatous Optic Disc Phenotypes. *Invest Ophthalmol Vis Sci*. 2015;56(10):5777-84.

Mayama C, Saito H, Hirasawa H, Tomidokoro A, Araie M, Iwase A, Ohkubo S, Sugiyama K, Hangai M, Yoshimura N. Diagnosis of early-stage glaucoma by grid-wise macular inner retinal layer thickness measurement and effect of compensation of disc-fovea inclination. *Invest Ophthalmol Vis Sci*. 2015;56(9):5681-90.

Kawashima Y, Uji A, Ooto S, Hangai M, Hosoda Y, Yoshimura N. Association between insufficient photoreceptor layer plugging and postoperative visual outcome in the surgically closed macular hole. *Am J Ophthalmol*. 2015;160(5):982-989.

Suda K, Hangai M, Akagi T, Noma H, Kimura Y, Hasegawa T, Yamada H, Yoshikawa M, Nakanishi H, Ikeda HO, Yoshimura N. Comparison of longitudinal changes in functional and structural measures for evaluating progression of glaucomatous optic neuropathy. *Invest Ophthalmol Vis Sci*. 2015;56(9):5477-84.

Kumagai K, Hangai M, Ogino N, Larson E. Effect of internal limiting membrane peeling on long-term visual outcomes for diabetic macular edema. *Retina*. 2015;35(7):1422-8.

Nakanishi H, Akagi T, Hangai M, Kimura Y, Suda K, Kumagai KK, Morooka S, Ikeda HO, Yoshimura N. Sensitivity and specificity for detecting early glaucoma in eyes with high myopia from normative database of macular ganglion cell complex thickness obtained from normal non-myopic or highly myopic Asian eyes. *Graefes Arch Clin Exp Ophthalmol*. 2015;253(7):1143-52.

Ooto S, Hangai M, Yoshimura N. Effects of sex and age on the normal retinal and choroidal structures on optical coherence tomography. *Curr Eye Res*. 2015;40(2):213-25.

Shoji T, Kuroda H, Suzuki M, Baba M, Hangai M, Araie M, Yoneya S. Correlation between lamina cribrosa tilt angles, myopia and glaucoma using OCT with a wide bandwidth femtosecond mode-locked laser. *PLoS One*. 2014;9(12): e116305.

Kimura Y, Akagi T, Hangai M, Takayama K, Hasegawa T, Suda K, Yoshikawa M, Yamada H, Nakanishi H, Unoki N, Ikeda HO, Yoshimura N. Lamina cribrosa defects and optic disc morphology in primary open angle glaucoma with high myopia. *PLoS One*. 2014;9(12):e115313.

Miyake M, Yamashiro K, Akagi-Kurashige Y, Oishi A, Tsujikawa A, Hangai M, Yoshimura N. Analysis of fundus shape in highly myopic eyes by using curvature maps constructed from optical coherence tomography. *PLoS One*. 2014;9(9):e107923.

Hosoda Y, Uji A, Hangai M, Morooka S, Nishijima K, Yoshimura N. Relationship between Retinal Lesions and Inward Choroidal Bulging in Vogt-Koyanagi-Harada Disease. *Am J Ophthalmol*. 2014;157(5):1056-63.

Hangai M, Ikeda HO, Akagi T, Yoshimura N. Paracentral scotoma in glaucoma detected by 10-2 but not by 24-2 perimetry. *Jpn J Ophthalmol*, in press.

Yoshikawa M, Akagi T, Hangai M, Ohashi-Ikeda H, Takayama K, Morooka S, Kimura Y, Nakano N, Yoshimura N. Alterations in the neural and connective tissue components of glaucomatous cupping after glaucoma surgery using swept-source optical coherence

tomography. *Invest Ophthalmol Vis Sci.* 2014;55:477-84.

Yamada H, Hangai M, Nakano N, Takayama K, Kimura Y, Miyake M, Akagi T, Ikeda HO, Noma H, Yoshimura N. Asymmetry analysis of macular inner retinal layers for glaucoma diagnosis. *Am J Ophthalmol.* 2014 Dec;158(6):1318-1329.

High-resolution imaging of photoreceptors in macular microholes. Ooto S, Hangai M, Takayama K, Ueda-Arakawa N, Makiyama Y, Hanebuchi M, Yoshimura N. *Invest Ophthalmol Vis Sci.* 2014 Aug 21;55(9):5932-43.

Ohkubo S, Higashide T, Udagawa S, Sugiyama K, Hangai M, Yoshimura N, Mayama C, Tomidokoro A, Araie M, Iwase A, Fujimura T. Focal relationship between structure and function within the central 10 degrees in glaucoma. *Invest Ophthalmol Vis Sci.* 2014;55(8):5269-77.

Yoshikawa M, Akagi T, Hangai M, Ohashi-Ikeda H, Takayama K, Morooka S, Kimura Y, Nakano N, Yoshimura N. Alterations in the neural and connective tissue components of glaucomatous cupping after glaucoma surgery using swept-source optical coherence tomography. *Invest Ophthalmol Vis Sci.* 2014;55(1):477-84.

Cone abnormalities in fundus albipunctatus associated with RDH5 mutations assessed using adaptive optics scanning laser ophthalmoscopy. Makiyama Y, Ooto S, Hangai M, Ogino K, Gotoh N, Oishi A, Yoshimura N. *Am J Ophthalmol.* 2014;157(3):558-70.

Nukada M, Hangai M, Mori S, Takayama K, Nakano N, Morooka S, Ikeda HO, Akagi T, Nonaka A, Yoshimura N. Imaging of localized retinal nerve fiber layer defects in preperimetric glaucoma using spectral-domain optical coherence tomography. *J Glaucoma.* 2014;23(3):150-9.

Uji A, Ooto S, Hangai M, Arichika S, Yoshimura N. Image quality improvement in adaptive optics scanning laser ophthalmoscopy assisted capillary visualization using B-spline-based elastic image registration. *PLoS One.* 2013;8:e80106

Makiyama Y, Ooto S, Hangai M, Takayama K, Uji A, Oishi A, Ogino K, Nakagawa S, Yoshimura N. Macular cone abnormalities in retinitis pigmentosa with preserved central vision using adaptive optics scanning laser ophthalmoscopy. *PLoS One.*

2013;8:e79447

Makiyama Y, Ooto S, Hangai M, Ogino K, Gotoh N, Oishi A, Yoshimura N. Cone Abnormalities in Fundus Albipunctatus Associated With RDH5 Mutations Assessed Using Adaptive Optics Scanning Laser Ophthalmoscopy. *Am J Ophthalmol*. 2014;157:558-570.e4.

Kimura Y, Hangai M, Matsumoto A, Akagi T, Ikeda HO, Ohkubo S, Sugiyama K, Iwase A, Araie M, Yoshimura N. Macular Structure Parameters as an Automated Indicator of Paracentral Scotoma in Early Glaucoma. *Am J Ophthalmol* 2013;156:907-917.e1

Mayama C, Saito H, Hirasawa H, Konno S, Tomidokoro A, Araie M, Iwase A, Ohkubo S, Sugiyama K, Otani T, Kishi S, Matsushita K, Maeda N, Hangai M, Yoshimura N. Circle- and grid-wise analyses of peripapillary nerve fiber layers by spectral domain optical coherence tomography in early-stage glaucoma. *Invest Ophthalmol Vis Sci* 2013;54:4519-26

Yokota S, Ooto S, Hangai M, Takayama K, Ueda-Arakawa N, Yoshihara Y, Hanebuchi M, Yoshimura N. Objective assessment of foveal cone loss ratio in surgically closed macular holes using adaptive optics scanning laser ophthalmoscopy. *PLoS One*. 2013;8:e63786.

Takayama K, Hangai M, Kimura Y, Morooka S, Nukada M, Akagi T, Ikeda HO, Matsumoto A, Yoshimura N. Three dimensional imaging of the lamina cribrosa defects in glaucoma using swept-source optical coherence tomography. *Invest Ophthalmol Vis Sci* 2013;54:4798-807.

Arichika S, Uji A, Hangai M, Ooto S, Yoshimura N. Noninvasive and direct monitoring of erythrocyte aggregates in human retinal microvasculature using Adaptive Optics Scanning Laser Ophthalmoscopy. *Invest Ophthalmol Vis Sci* 2013;54:4394-402.

Nakano N, Hangai M, Noma H, Nukada M, Mori S, Morooka S, Takayama K, Kimura Y, Ikeda HO, Akagi T, Yoshimura N. Macular imaging in highly myopic eyes with and without glaucoma. *Am J Ophthalmol* 2013;156:511-523.e6.

Kumagai K, Hangai M, Larson E, Ogino N. Progressive changes of regional macular thickness after macular hole surgery with internal limiting membrane peeling. *Invest*

Ophthalmol Vis Sci 2013;7:751-6.

Lee J, Hangai M, Kimura Y, Takayama K, Kee C, Yoshimura N. Measurement of macular ganglion cell layer and circumpapillary retinal nerve fiber layer to detect paracentral scotoma in early glaucoma. *Graefes Arch Clin Exp Ophthalmol* 2013;251:2003-12.

Kumagai K, Hangai M, Larson E, Ogino N. Foveal thickness in healthy fellow eyes of patients with unilateral macular holes. *Am J Ophthalmol* 2013;156:140-8.

Ooto S, Hangai M, Takayama K, Ueda-Arakawa N, Tsujikawa A, Yamashiro K, Oishi A, Hanebuchi M, Yoshimura N. Comparison of cone pathologic changes in idiopathic macular telangiectasia types 1 and 2 using adaptive optics scanning laser ophthalmoscopy. *Am J Ophthalmol* 2013;155:1045-1057.e4

Iida Y, Hangai M, Yoshikawa M, Ooto S, Yoshimura N. Local biometric features and visual prognosis after surgery for treatment of myopic foveoschisis. *Retina*. 2013;33:1179-87.

Kumagai K, Hangai M, Furukawa M, Larson E, Ogino N. Three-dimensional foveal shape changes after asymptomatic macular posterior vitreous detachment. *Clin Ophthalmol*. 2013;7:751-6

Wakazono T, Ooto S, Hangai M, Yoshimura N. Photoreceptor outer segment abnormalities and retinal sensitivity in acute zonal occult outer retinopathy. *Retina*. 2013;33(3):642-8.

Hirasawa H, Araie M, Tomidokoro A, Saito H, Iwase A, Ohkubo S, Sugiyama K, Ootani T, Kishi S, Matsushita K, Maeda N, Hangai M, Yoshimura N. Reproducibility of thickness measurements of macular inner retinal layers using SD-OCT with or without correction of ocular rotation. *Invest Ophthalmol Vis Sci*. 2013;54(4):2562-70

Nukada K, Hangai M, Ooto S, Yoshikawa M, Yoshimura N. Tomographic features of macula after successful macular hole surgery. *Invest Ophthalmol Vis Sci*. 2013;54(4):2417-28

Akagi T, Hangai M, Kimura Y, Ikeda HO, Nonaka A, Matsumoto A, Akiba M, Yoshimura N. Peripapillary scleral deformation and retinal nerve fiber damage in high myopia assessed with swept-source optical coherence tomography. *Am J Ophthalmol.* 2013;155:927-936.e1

Takayama K, Ooto S, Hangai M, Ueda-Arakawa N, Yoshida S, Akagi T, Ikeda HO, Nonaka A, Hanebuchi M, Inoue T, Yoshimura N. High-resolution imaging of retinal nerve fiber bundles in glaucoma using adaptive optics scanning laser ophthalmoscopy. *Am J Ophthalmol.* 2013;155:870-881.e3

Hirashima T, Hangai M, Nukada M, Nakano N, Morooka S, Akagi T, Nonaka A, Yoshimura N. Frequency-doubling technology and retinal measurements with spectral-domain optical coherence tomography in preperimetric glaucoma. *Graefes Arch Clin Exp Ophthalmol.* 2013;251:129-37.

Suehira N, Sotaro Ooto S, Hangai M, Matsumoto K, Tomatsu N, Yuasa T, Yamada K, Yoshimura N. Three-beam spectral-domain optical coherence tomography for retinal imaging. *J Biomed Opt.* 2012;17:106001.

Oishi M, Nakamura H, Hangai M, Oishi A, Otani A, Yoshimura N. Contrast visual acuity in patients with retinitis pigmentosa assessed by a contrast sensitivity tester. *Indian J Ophthalmol.* 2012;60:545-9.

Takayama K, Hangai M, Durbin M, Nakano N, Morooka S, Akagi T, Ikeda HO, Yoshimura N. A novel method to detect local ganglion cell loss in early glaucoma using spectral-domain optical coherence tomography. *Invest Ophthalmol Vis Sci.* 2012;53:6904-13.

Kimura Y, Hangai M, Morooka S, Takayama K, Nakano N, Nukada M, Ikeda HO, Akagi T, Yoshimura N. Retinal nerve fiber layer defects in highly myopic eyes with early glaucoma. *Invest Ophthalmol Vis Sci.* 2012;53:6472-8.

Furukawa H, Arimoto H, Shirai T, Ooto S, Hangai M, Yoshimura N. Oximetry of retinal capillaries by multicomponent analysis. *Appl Spectrosc.* 2012;66:962-9.

Morooka S, Hangai M, Nukada M, Nakano N, Takayama K, Kimura Y, Akagi T, Ikeda

HO, Nonaka A, Yoshimura N. Wide 3-dimensional macular ganglion cell complex imaging with spectral-domain optical coherence tomography in glaucoma. *Invest Ophthalmol Vis Sci.* 2012;53:4805-12.

Akagi T, Hangai M, Takayama K, Nonaka A, Ooto S, Yoshimura N. In vivo imaging of lamina cribrosa pores by adaptive optics scanning laser ophthalmoscopy. *Invest Ophthalmol Vis Sci.* 2012;53:4111-9.

Ellabban AA, Hangai M, Yamashiro K, Nakagawa S, Tsujikawa A, Yoshimura N. Tomographic fundus features in pseudoxanthoma elasticum: comparison with neovascular age-related macular degeneration in japanese patients. *Eye.* 2012;26:1086-94.

Kumagai K, Ogino N, Furukawa M, Hangai M, Kazama S, Nishigaki S, Larson E. Retinal thickness after vitrectomy and internal limiting membrane peeling for macular hole and epiretinal membrane. *Clin Ophthalmol.* 2012;6:679-88.

Jirarattanasopa P, Ooto S, Tsujikawa A, Yamashiro K, Hangai M, Hirata M, Matsumoto A, Yoshimura N. Assessment of macular choroidal thickness by optical coherence tomography and angiographic changes in central serous chorioretinopathy. *Ophthalmology.* 2012;119:1666-78.

Iwama D, Hangai M, Ooto S, Sakamoto A, Nakanishi H, Fujimura T, Domalpally A, Danis RP, Yoshimura N. Automated assessment of drusen using three-dimensional spectral-domain optical coherence tomography. *Invest Ophthalmol Vis Sci* 2012;53:1576-83.

Uji A, Hangai M, Ooto S, Takayama K, Arakawa N, Imamura H, Nozato K, Yoshimura N. The source of moving particles in parafoveal capillaries detected by adaptive optics scanning laser ophthalmoscopy. *Invest Ophthalmol Vis Sci.* 2012;53:171-8.

Kumagai K, Ogino N, Hangai M, Larson E. Percentage of fellow eyes that develop full-thickness macular hole in patients with unilateral macular hole. *Arch Ophthalmol* 2012;130:393-4.

Ellabban AA, Tsujikawa A, Matsumoto A, Ogino K, Hangai M, Ooto S, Yamashiro K, Akiba M, Yoshimura N. Macular choroidal thickness and volume in eyes with angiod

streaks measured by swept-source optical coherence tomography. *Am J Ophthalmol* 2012;153:1133-1143.

Takayama K, Ooto S, Hangai M, Arakawa N, Oshima S, Hanebuchi M, Inoue T, Yoshimura N. High-resolution imaging of the retinal nerve fiber layer in normal eyes using adaptive optics scanning laser ophthalmoscopy. *PLoS ONE* 2012;7:e33158.

Ooto S, Hangai M, Takayama K, Ueda-Arakawa N, Hanebuchi M, Yoshimura N. Photoreceptor damage and foveal sensitivity in surgically closed macular holes: an adaptive optics scanning laser ophthalmoscopy study. *Am J Ophthalmol*. 2012;154:174-186.

Muraoka Y, Ikeda HO, Nakano N, Hangai M, Toda Y, Okamoto-Furuta K, Kohda H, Kondo M, Terasaki H, Kakizuka A, Yoshimura N. Real-time imaging of rabbit retina with retinal degeneration by using spectral-domain optical coherence tomography. *PLoS One*. 2012;7(4):e36135.

Suda K, Hangai M, Yoshimura N. Axial length and outcomes of macular hole surgery assessed by spectral-domain optical coherence tomography. *Am J Ophthalmol* 2011;151:118-127.e1.

Ooto S, Hangai M, Takayama K, Sakamoto A, Tsujikawa A, Oshima S, Inoue T, Yoshimura N. High-resolution imaging of the photoreceptor layer in epiretinal membrane using adaptive optics scanning laser ophthalmoscopy. *Ophthalmology* 2010;117:1800-9.

Nakamura H, Hangai M, Mori S, Hirose F, Yoshimura N. Hemispherical focal macular photopic negative response and macular inner retinal thickness in open-angle glaucoma. *Am J Ophthalmol* 2011;151:494-506.e1.

Nukada M, Hangai M, Mori S, Nakano N, Nakanishi H, Ohashi-Ikeda H, Nonaka A, MD, Yoshimura N. Detection of localized retinal nerve fiber layer defects in glaucoma using enhanced spectral domain optical coherence tomography. *Ophthalmology* 2011;118:1038-48

Kotera Y, Hangai M, Hirose F, Mori S, Yoshimura N. Three-dimensional imaging of

macular inner structures in glaucoma using spectral-domain optical coherence tomography. *Invest Ophthalmol Vis Sci* 2011;52:1412-21.

Kumagai K, Hangai M, Larson E, Ogino N. Vitreoretinal interface and foveal deformation in asymptomatic fellow eyes of patients with unilateral macular holes. *Ophthalmology* 2011;118:1638-44.

Hirata M, Tsujikawa A, Matsumoto A, Hangai M, Ooto S, Yamashiro K, Akiba M, Yoshimura N. Macular choroidal thickness and volume in normal subjects measured by swept-source optical coherence tomography. *Invest Ophthalmol Vis Sci*. 2011;52:4971-8.

Nonaka A, Hangai M, Akagi T, Mori S, Nukada M, Nakano N, Yoshimura N. Biometric features of peripapillary atrophy Beta in eyes with high myopia. *Invest Ophthalmol Vis Sci*. 2011;52:6706-13.

Ooto S, Hangai M, Takayama K, Arakawa N, Tsujikawa A, Koizumi H, Oshima S, Yoshimura N. High-resolution photoreceptor imaging in idiopathic macular telangiectasia type 2 using adaptive optics scanning laser ophthalmoscopy. *Invest Ophthalmol Vis Sci*. 2011;52:5541-50.

Nakano N, Hangai M, Nakanishi H, Mori S, Nukada M, Kotera Y, Ikeda OH, Nakamura H, Nonaka A, Yoshimura N. Macular ganglion cell layer imaging in preperimetric glaucoma with speckle-noise-reduced spectral-domain optical coherence tomography. *Ophthalmology* 2011;118:2414-26.

Ooto S, Hangai M, Tomidokoro A, Saito H, Araie M, Otani T, Kishi S, Matsushita K, Maeda N, Shirakashi M, Abe H, Ohkubo S, Sugiyama K, Iwase A, Yoshimura N. Effects of age, sex, and axial length on the three-dimensional profile of normal macular layer structures. *Invest Ophthalmol Vis Sci* 52: 8769-8779, 2011. Nakano N, Ikeda HO, Hangai M, Muraoka Y, Toda Y, Kakizuka A, Yoshimura N. Longitudinal and simultaneous imaging of retinal ganglion cells and inner retinal layers in a mouse model of glaucoma induced by N-methyl-D-aspartate. *Invest Ophthalmol Vis Sci* 2011;52:8754-8762. Haruta M, Hangai M, Taguchi C, Yamakawa R. Spectral-domain optical coherence tomography of the choroid in choroidal osteoma. *Ophthalmic Surg Lasers Imaging* 2011;42:118-121. Ooto S, Hangai M, Yoshimura N. Photoreceptor restoration in unilateral acute idiopathic maculopathy on adaptive optics scanning laser ophthalmoscopy. *Arch Ophthalmol* 129:1633-1635, 2011. Ooto S, Hangai M, Sakamoto A,

Tomidokoro A, Araie M, Otani T, Kishi S, Matsushita K, Maeda N, Shirakashi M, Abe H, Takeda H, Sugiyama K, Saito H, Iwase A, Yoshimura N. Three-dimensional profile of macular retinal thickness in normal Japanese eyes. *Invest Ophthalmol Vis Sci*. 2010;51:465-73. Arichika S, Hangai M, Yoshimura N. Correlation between thickening of the inner and outer retina and visual acuity in patients with epiretinal membrane. *Retina* 30:503-508:2010. Nakano N, Hangai M, Nakanishi H, Inoue R, Unoki N, Hirose F, Ojima T, Yoshimura N. Bleb, GreafeEarly trabeculectomy bleb walls on anterior-segment optical coherence tomography. *Graefes Arch Clin Exp Ophthalmol*. 2010;248:1173-82 Mori S, Hangai M, Sakamoto A, Yoshimura N. Spectral-domain Optical Coherence Tomography Measurement of Macular Volume for Diagnosing Glaucoma. *J Glaucoma* 2010;19:528-34. Sakamoto A, Hangai M, Nukada M, Nakanishi H, Mori S, Kotera Y, Inoue R, Yoshimura N. Three-Dimensional Imaging of Macular Retinal Nerve Fiber Layer in Glaucoma Using Spectral-Domain Optical Coherence Tomography. *Invest Ophthalmol Vis Sci* 2010;51:5062-70. Yang Q, Reisman CA, Wang Z, Fukuma Y, Hangai M, Yoshimura N, Tomidokoro A, Araie M, Raza AS, Hood DC, Chan A. Automated layer segmentation of macular OCT images using dual-scale gradient information. *Optics Express*, 2010;18:21293-307. Ooto S, Hangai M, Sakamoto A, Tsujikawa A, Yamashiro K, Ojima Y, Yamada Y, Mukai H, Oshima S, Inoue T, Yoshimura N. High-resolution imaging of resolved central serous chorioretinopathy using adaptive optics scanning laser ophthalmoscopy. *Ophthalmology* 2010;117:1800-9. Hirasawa H, Tomidokoro A, Araie M, Konno S, Maeda N, Sugiyama K, Iwase A, Kishi S, Abe H, Hangai M, Yoshimura N. Three-dimensional profile of peripapillary retinal nerve fiber layer thickness and its related factors in normal eyes. *Arch Ophthalmol* 2010;128:1420-6. Fujimoto M, Hangai M, Suda K, Yoshimura N. Features associated with foveal retinal detachment in myopic macular retinoschisis. *Am J Ophthalmol*. 2010 Dec;150(6):863-70. Kameda T, Tanabe T, Hangai M, Ojima T, Aikawa H, Yoshimura N. Fixation behavior in advanced stage glaucoma assessed by the MicroPerimeter MP-1. *Jpn J Ophthalmol*. 2009;53:580-7. Inoue R, Hangai M, Kotera Y, Nakanishi H, Mori S, Morishita S, Yoshimura N. Three-dimensional high-speed optical coherence tomography imaging of lamina cribrosa in glaucoma. *Ophthalmology* 116: 214-222, 2009. Ojima Y, Hangai M, Sakamoto A, Tsujikawa A, Otani A, Tamura H, Yoshimura N. Improved visualization of polypoidal choroidal vasculopathy lesions using spectral-domain optical coherence tomography. *Retina* 29: 52-59, 2009. Nakanishi H, Hangai M, Unoki N, Sakamoto A, Tsujikawa A, Kita M, Yoshimura N. Spectral-domain optical coherence tomography imaging of the detached macula in rhegmatogenous retinal detachment. *Retina* 29: 232-242, 2009. Hangai M, Fujimoto M, Yoshimura N. Morphology and

function in multiple evanescent white dot syndrome. *Arch Ophthalmol*, 2009;127:1307-13. Kotera Y, Yasuno Y, Hangai M, Inoue R, Makita S, Nakanishi H, Yamanari M, Yoshimura N. Comparison of spectral domain optical coherence tomography and color photographic imaging of the optic nerve head in management of glaucoma. *Ophthalmic Surg Lasers Imaging*. 2009;40:255-63. Ishihara K, Hangai M, Kita M, Yoshimura N. Acute Vogt-Koyanagi-Harada disease in enhanced spectral-domain optical coherence tomography. *Ophthalmology*. 2009;116:1799-807. Hangai M, Yamamoto M, Sakamoto A, Yoshimura N. Ultrahigh-resolution versus speckle noise-reduction in spectral-domain optical coherence tomography. *Optics Express* 17: 4221-4235, 2009. Morishita S, Tanabe T, Yu S, Hangai M, Ojima T, Aikawa H, Yoshimura N. Retinal nerve fibre layer assessment in myopic glaucomatous eyes: comparison of GDx variable corneal compensation with GDx enhanced corneal compensation. *Br J Ophthalmol* 92: 1377-1381, 2008. Ojima Y, Tsujikawa A, Hangai M, Nakanishi H, Inoue R, Sakamoto A, Yoshimura N. Retinal sensitivity measured with the micro perimeter 1 after resolution of central serous chorioretinopathy. *Am J Ophthalmol* 146: 77-84, 2008. Sakamoto A, Hangai M, Yoshimura N. Spectral-domain optical coherence tomography with multiple B-scan averaging for enhanced imaging of retinal diseases. *Ophthalmology* 115: 1071-1078: 2008. Yamaike N, Tsujikawa A, Ota M, Sakamoto A, Kotera Y, Kita M, Miyamoto K, Yoshimura N, Hangai M. Three-dimensional Imaging of cystoid macular edema in retinal vein occlusion. *Ophthalmology* 115: 355-362, 2008. Ojima Y, Hangai M, Sasahara M, Gotoh N, Inoue R, Yasuno Y, Makita S, Yatagai T, Tsujikawa A, Yoshimura N. Three-dimensional imaging of the foveal photoreceptor layer in central serous chorioretinopathy using high-speed optical coherence tomography. *Ophthalmology* 114: 2197-2207, 2007. Hangai M, Ojima Y, Gotoh N, Inoue R, Yasuno Y, Makita S, Yamanari M, Yatagai T, Kita M, Yoshimura N. Three dimensional imaging of macular holes with high-speed optical coherence tomography. *Ophthalmology* 114: 763-773, 2007. Unoki N, Nishijima K, Sakamoto A, Kita M, Watanabe D, Hangai M, Kimura T, Kawagoe N, Ohta M, Yoshimura N. Retinal sensitivity loss and structural disturbance in areas of capillary nonperfusion of eyes with diabetic retinopathy: *Am J Ophthalmol* 144: 755-760, 2007. Ojima T, Tanabe T, Hangai M, Yu S, Morishita S, Yoshimura N. Measurement of retinal nerve fiber layer thickness and macular volume for glaucoma detection using optical coherence tomography. *Jpn J Ophthalmol* 51: 197-203, 2007. Yu S, Tanabe T, Hangai M, Morishita S, Kurimoto Y, Yoshimura N. Scanning laser polarimetry with variable corneal compensation and optical coherence tomography in tilted disk. *Am J Ophthalmol* 142 :475-482, 2006. Tsujikawa A, Hangai M, Kikuchi M, Ishida K, Kurimoto Y. Visual

field defect after radial optic neurotomy for central retinal vein occlusion. *Jpn J Ophthalmol* 50: 158-160, 2006. Hangai M, He S, Hoffmann S, Lim JI, Ryan SJ, Hinton DR. Sequential induction of angiogenic growth factors by TNF- α in choroidal endothelial cells. *J Neuroimmunol* 171: 45-56, 2006. Lim JI, Spee C, Hangai M, Rocha J, Ying HS, Ryan SJ, Hinton DR. Neuropilin-1 expression by endothelial cells and retinal pigment epithelial cells in choroidal neovascular membranes. *Am J Ophthalmol* 140: 1044-1050, 2005. Yamamoto K, Tsujikawa A, Hangai M, Fujihara M, Iwawaki T, Kurimoto Y. Neovascular glaucoma after branch retinal artery occlusion. *Jpn J Ophthalmol* 49: 388-390, 2005. Hangai M, Kitaya N, Xu J, Chan CK, Kim JJ, Werb Z, Ryan SJ, Brooks PC. Matrix metalloproteinase-9-dependent exposure of a cryptic migratory control site in collagen is required before retinal angiogenesis. *Am J Pathol* 161: 1429-1437, 2002. Xu J, Rodriguez D, Petitclerc E, Kim JJ, Hangai M, Moon YS, Davis GE, Brooks PC. Proteolytic exposure of a cryptic site within collagen type IV is required for angiogenesis and tumor growth in vivo. *J Cell Biol* 154: 1069-1079, 2001. Hangai M, Moon YS, Kitaya N, Chan CK, Wu DY, Peters KG, Ryan SJ, Hinton DR. Systemically expressed soluble Tie2 inhibits intraocular neovascularization. *Hum Gene Ther* 12: 1311-1321, 2001. Hangai M, Murata T, Miyawaki N, Spee C, Lim JI, He S, Hinton DR, Ryan SJ. Angiopoietin-1 upregulation by vascular endothelial growth factor in human retinal pigment epithelial cells. *Invest Ophthalmol Vis Sci* 42: 1617-1625, 2001. Murata T, He S, Hangai M, Ishibashi T, Xi XP, Kim S, Hsueh WA, Ryan SJ, Law RE, Hinton DR. Peroxisome proliferator-activated receptor-gamma ligands inhibit choroidal neovascularization. *Invest Ophthalmol Vis Sci* 41: 2309-2317, 2000. Inatani M, Tanihara H, Honjo M, Hangai M, Kresse H, Honda Y. Expression of Proteoglycan Decorin in Neural Retina *Invest Ophthalmol Vis Sci* 40: 1783-1791, 1999. Akimoto M, Miyatake S, Okazaki K, Kogishi J, Hangai M, Honda Y. Adenovirally expressed basic fibroblast growth factor rescues photoreceptor cells in RCS rats. *Invest Ophthalmol Vis Sci* 40: 273-279, 1999. Kogish J, Miyatake S, Hangai M, Akimoto M, Okazaki K, Honda Y. Mutant herpes simplex virus-mediated suppression of retinoblastoma. *Curr Eye Res* 18: 321-326, 1999.

Hangai M, Miyamoto K, Hiroi K, Tsujikawa A, Ogura Y, Honda Y, Yoshimura N. Roles of constitutive nitric oxide synthase in postischemic rat retina. *Invest Ophthalmol Vis Sci* 40: 450-458, 1999. Hangai M, Yoshimura N, Hiroi K, Mandai M, Honda Y. Role of nitric oxide during the initial phase of reperfusion following retinal ischemia in the rat. *Ophthalmic Res* 31: 16-23, 1999. Miyatake SI, Tani S, Feigenbaum F, Sundaresan P, Toda H, Narumi O, Kikuchi H, Hashimoto N, Hangai M, Martuza RL, Rabkin SD. Hepatoma-specific anti-tumor activity of an albumin enhancer/promoter regulated

herpes simplex virus in vivo. *Gene Therapy* 6: 564-572, 1999. Akimoto M, Hangai M, Okazaki K, Kogishi J, Honda Y, Kaneda Y. Growth inhibition of cultured human Tenon's fibroblastic cells by targeting the E2F transcription factor. *Exp Eye Res* 67: 395-401, 1998. Murata T, Hangai M, Spee C, Gordon EM, Anderson WF, Hinton DR, Ryan SJ. Retrovirus-mediated gene transfer to photocoagulation-induced choroidal neovascular membrane. *Invest Ophthalmol Vis Sci* 39: 2474-2478, 1998. Ishibashi T, Murata T, Hangai M, Nagai R, Seikoh H, Lopez PF, Hinton DR, Ryan SJ. Advanced glycation end products in age-related macular degeneration. *Arch Ophthalmol* 116: 1629-1632, 1998. Hangai M, Tanihara H, Honda Y, Kaneda Y. In vivo delivery of phosphorothioate oligonucleotides into murine retina. *Arch Ophthalmol* 116: 342-348, 1998. Hangai M, Tanihara H, Honda Y, Kaneda Y. Introduction of DNA into the rat and primate trabecular meshwork by fusogenic liposomes. *Invest Ophthalmol Vis Sci* 39: 509-516, 1998. Honjo M, Tanihara H, Negi A, Hangai M, Taniguchi T, Matsumura M, Nagata M. Trabeculectomy ab externo, phacoemulsification and implantation of an intraocular lens: Preliminary report of surgical results and procedures. *J Cataract Refract Surg* 22: 601-606, 1996. Hangai M, Kaneda Y, Tanihara H, Honda Y. In vivo gene transfer into retina mediated by a novel liposome system. *Invest Ophthalmol Vis Sci* 37: 2678-2685, 1996. Mandai M, Mittag TW, Kogishi J, Iwaki M, Hangai M, Yoshimura N. Roles of nitric oxide synthase isozymes in endotoxin induced uveitis. *Invest Ophthalmol Vis Sci* 37: 826-832, 1996. Hangai M, Yoshimura N, Honda Y. Increased cytokine gene expression in rat retina following transient ischemia. *Ophthalmic Res* 28: 248-254, 1996. Hangai M, Yoshimura N, Hiroi K, Mandai M, Honda Y. Inducible nitric oxide synthase in retinal ischemia-reperfusion injury. *Exp Eye Res* 63: 501-509, 1996. Matsuda T, Tanihara H, Hangai M, Chihara E. Surgical results and complications of trabeculectomy with intraoperative application of mitomycin C. *Jpn J Ophthalmol* 40: 526-532, 1996. Tanihara H, Hangai M, Sawaguchi S, Abe H, Kageyama M, Nakazawa F, Shirasawa E, Honda Y. Up-regulation of glial fibrillary acidic protein in the retina of eyes with experimental glaucoma. *Arch Ophthalmol* 115: 752-756, 1997. Chihara E, Liu X, Dong J, Takashima Y, Akimoto M, Hangai M, Kuriyama S, Tanihara H, Hosoda M, Tsukahara S. Severe myopia as a risk factor for progressive visual field loss in eyes with treated primary open angle glaucoma. *Ophthalmologica* 211: 66-71, 1997. Yoshida M, Yoshimura N, Hangai M, Tanihara H, Honda Y. Interleukin-1 α , interleukin-1 β , and tumor necrosis factor gene expression in endotoxin-induced uveitis. *Invest Ophthalmol Vis Sci* 35: 1107-1113, 1994. Hangai M, Yoshimura N, Yoshida M, Yabuuchi K, Honda Y. Interleukin-1 gene expression in transient retinal ischemia in the rat. *Invest Ophthalmol Vis Sci* 36: 571-578, 1995.